




BASS

British Association of Spine Surgeons

Lumbar Spondylolysis and Spondylitic Spondylolisthesis – Surgical Options




Issue 4: September 2021
Review date: August 2024

Following your recent investigations and consultation with your spinal surgeon, it has been confirmed that your symptoms are caused by **lumbar spondylolysis**.

This is a stress fracture of the narrow bridge of bone between the facet joints (*pars interarticularis*) at the back of the spine, commonly called a **pars defect**. There may be a hereditary aspect to spondylolysis, for example an individual may be born with thin vertebral bone and therefore be vulnerable to this condition; **or** certain sports, such as gymnastics, weight lifting and football can put a great deal of stress on the bones through constantly over-stretching the spine. Either can result in a stress fracture on one or both sides of the vertebra (bone of the spine).

Many people are not aware of their stress fracture or experience any problems but symptoms can occur, including lower back pain, pain in the thighs and buttocks, stiffness, muscle tightness and tenderness. The diagnosis is often confirmed with an injection of an anti-inflammatory drug (steroid) into the *pars defect*, when it is followed by successful relief from pain (even if only temporary).





If the stress fracture weakens the bone so much that it is unable to maintain its proper position, the vertebra can start to shift out of place. This condition is called **spondylolisthesis**.

The normal spinal column has a central canal (or passage) through which the spinal nerves pass down. At each side of the vertebral level, spinal nerve roots branch out to each side. The solid spinal cord stops at the top of the lumbar spine (lower back) and from this point the nerves to the bottom and legs pass through the lower canal like a horse's tail forming the cauda equina. The spinal cord, nerve roots and cauda equina are surrounded by cerebrospinal fluid (CSF) and are all contained within a membrane, or covering, called the dura mater, rather like the thin layer that covers a boiled egg.

In spondylolisthesis there is a forward slippage of one lumbar vertebra on the vertebra below it. The degree of spondylolisthesis may vary from mild to severe but if too much slippage occurs, the nerve roots can be stretched where they branch out of the spinal canal. This can cause nerve symptoms.

X-ray showing spondylolisthesis at L5/S1



A nerve is like an electrical wire. It tells your muscles to move and gives your brain information about various sensations, such as pain, temperature, light touch, pressure sensation and the position of your legs. Lumbar nerve root pain or radiculopathy (often called sciatica) generally goes below the knee and is felt in the area of the leg that the spinal nerve supplies. Symptoms also associated with sciatica include altered sensation, pins and needles, burning, numbness or even weakness of the muscles in the leg the nerve supplies.

Treatment varies depending on the severity of the condition. Most patients only require treatment such as physiotherapy, medication and manipulative therapy, combined with some lifestyle changes, like avoiding contact sports, weight lifting and leaning too far backwards (hyperextension).

For patients whose pain does not settle with treatment, surgery may be necessary. The actual procedure may differ depending on the severity of the condition. For patients who have no, or minimal slippage, surgery may simply involve directly repairing the pars defect with a screw and bone graft.

But for patients where slippage (spondylolisthesis) occurs resulting in nerve pain (sciatica), surgery needs to address both the problem with the nerves, as well as the instability of the spine. A decision then needs to be made whether an internal system of screws, rods and bone graft is needed to hold together the vertebra.

The nature of spinal surgery is not to 'cure' a condition but is aimed to provide benefit with a good percentage improvement and relief of symptoms. Good relief from back and leg symptoms following surgery usually occurs in approximately 70% of cases (up to seven out of 10 people). This is not necessarily felt immediately but over a period of time (often several months). Sometimes however, numbness or weakness can persist (even with a technically successful operation).


The operation

There are several different techniques when performing an operation to deal with the condition spondylolysis and spondylitic spondylolisthesis. Expected outcomes from all methods are very similar and the choice of operation will be decided by the surgeon. Sometimes, the exact extent of surgery can only be decided during the procedure.

Direct Pars repair

This is performed through an incision in the midline of the lower back. The muscles are held apart to gain access to the bony arch and roof of the spine (lamina). The fractured bone is identified and a screw passed directly across from one side to another, joining it together.

X-ray showing direct repair using screws, side and front views



Decompression

- 1 Laminectomy and/or laminotomy. A similar approach to the direct Pars repair is used. Next, the surgeon needs to gain entry into the spinal canal by removing some bone, either by cutting away the whole area of lamina (laminectomy) or making a small window in the lamina (laminotomy) with a high-speed burr (like a dentist's drill). Further bone and inflammatory tissue around the fracture site is removed and often the facet joints, which are directly over the nerve roots, are undercut (trimmed), to relieve the pressure on the nerves and give them a wider passage as they pass out of the spine.
- 2 Minimally invasive (tubular) decompression. With this approach, the surgeon attempts to reduce muscle dissection and injury by working through a narrow tube.


Stabilisation

- 1 Bone graft. This is used to fuse (join together) and stabilise the spine in conjunction with other techniques, which are mentioned next. It is laid between the outer segments of the spine, in between the transverse process (intertransverse region) and your new bone will, over time, grow into the bone graft. This is a biological process over 3–6 months, known as spinal fusion. There are several techniques to get bone graft needed for spinal fusion:
 - **patient's own bone (autograft bone).** The bone that is removed during surgery can be used as a bone graft. If more is needed, for the most part, **artificial bone** is now used, although it can be taken from the pelvis (iliac crest) if required;


- **donor bone (allograft bone).** Donor bone graft does not contain living bone cells but acts as calcium scaffolding which your own bone grows into and eventually replaces.

2 Pedicle screw fixation. This is a system of screws and rods which hold the vertebra together and prevents movement at the segment that is being fused, like an internal scaffolding system. Screws are placed into the part of the vertebra called the pedicle, which go directly from the back of the spine to the front, on both sides, above and below the unstable spinal segment. These screws then act as firm anchor points to which rods can be connected. After the bone graft grows and fuses to the spine (after many months), the rods and screws are no longer needed for stability. However, most surgeons do not recommend removing them except in rare cases.

Side view of pedicle screw placement




Overhead view of pedicle screw in vertebral body



Complete system in place

X-rays showing the stabilisation system in place, side and front views




3 Interbody fusion. This is a procedure where the surgeon will remove the intervertebral disc, the structure between the bones of the spine (vertebra) and fuse the space with a cage and bone graft. This can enable the surgeon to achieve an all round (360°) spinal fusion when used in conjunction with pedicle screw fixation and to increase the height of the interbody space, and therefore the bone canal, for the nerves to pass through. Before interbody fusion surgery is carried out, several factors are taken into consideration. These include how narrow the nerve canal has become and how much the vertebra has slipped.

4 Intervertebral fusion cages. This is like a hollow Lego brick which props up the disc space between the two bones (vertebra). It is a tight fit and gives immediate stability. The cage is different widths, heights and depths to fit your spine exactly. It is made from carbon fibre or titanium metal. They can be filled with bone graft or artificial bone if required and are used in interbody fusion surgery.



The surgical approach (way in) to interbody fusion also varies and can include:

- posterior lumbar interbody fusion (PLIF): The spine is approached from the back and part of the lamina bone and facet joint is removed on both sides to gain access to the disc.

Posterior lumbar interbody fusion (PLIF)





X-rays showing PLIF cages in place, side and front views





- transforaminal lumbar interbody fusion (TLIF): The spine is approached again from the back and all of the facet joint is removed on one side to gain access to the disc.

Transforaminal lumbar interbody fusion (TLIF)

facet joint removed to gain access to the disc for removal




X-rays showing TLIF cage in place, side and front views




- anterior lumbar interbody fusion (ALIF): The spine is approached through an incision in the tummy. The abdominal contents lay inside a large sack (peritoneum) that can be retracted (moved to the side), allowing the surgeon access to the front of the spine without actually entering the abdomen. The large blood vessels (aorta, common iliac artery and vena cava) that lie over the front of the spine are carefully moved aside. Sometimes surgery is performed in conjunction with a vascular surgeon, who will mobilise the blood vessels if there are any concerns or difficulty

Diagram of cage placement



Anterior lumbar interbody fusion (ALIF)



X-rays showing ALIF cage in place, side and front views



with this.

After the blood vessels have been moved aside, a 'window' is cut in the anterior ligament and fibrous wall of the disc (annulus fibrosus), retaining as much of it as possible to provide stability for the cage. The disc material (nucleus pulposus) is then removed and the cage, containing bone graft, is placed in the space created. Some cages require screwing into the vertebral body above and below them. Your own bone will, over time, grow into the bone graft.

This particular approach to the spine has risks and complications which are specific to it and are listed below the general ones.

Risks and complications

As with any form of surgery, there are risks and complications associated with this procedure.

These can include:

- damage to a nerve root. This occurs in less than 1 out of 100 cases of primary surgery but is much more common in revision or 're-do' surgeries where injury can occur in up to 10 out of 100 cases. If this happens, you may get weakness in the muscles supplied by that particular nerve root and/or numbness, tingling or hypersensitivity in the area of skin it supplies;
- tearing of the outer lining or covering which surrounds the nerve roots (dura). This is reported in fewer than 5 out of 100 cases. It may occur as a result of the bone being very stuck to the lining and tearing it as the bone is lifted off. Again, it is much more common in 're-do' surgery. Usually the hole or tear in the dura is repaired with stitches, a patch or a special glue. If the puncture or hole re-opens you may get cerebrospinal fluid (CSF) leaking from the wound, headaches or, very rarely, meningitis. Although rare, the problems of leakage can persist. This could result in you having to return to theatre to enable the surgeon to revise the repair of the dura but the risk of a second operation being required within a few days or weeks is less than 0.05%;
- recurrent leg pain as a result of scarring;
- problems with positioning during the operation which might include skin injuries or pressure problems. Special gel mattresses and operating tables are used to minimise this;
- infection. Superficial wound infections may occur in up to 4 out of 100 cases. These are often easily treated with a course of antibiotics. Deep wound infections may occur in fewer than 1 out of 100 cases. These can be more difficult to treat with antibiotics alone and sometimes patients require more surgery to clean out the infected tissue. This risk may increase for people who have diabetes, an impaired immune system or are taking steroids;
- blood clots (thromboses) in the deep veins of the legs (DVT) or lungs (PE). These occur when the blood in the large veins of the leg forms blood clots and may cause the leg to swell and become painful and warm to the touch. Although rare, if not treated this

could be a fatal condition if the blood clot travels from the legs to the lungs, cutting off the blood supply to a portion of the lung. It is reported as happening in fewer than 1 in 700 cases. There are many ways to reduce the risk of blood clots forming. The most effective is to get moving as soon as possible after your operation. Walk regularly as soon as you are able to, both in hospital and when you return home. Perform the leg exercises as shown to you by the physiotherapist and keep well hydrated by drinking plenty of water. Ladies are also advised to stop taking any medication which contains the hormone oestrogen (like the combined contraceptive pill or HRT) four weeks before surgery, as taking this during spinal surgery can increase the chances of developing a blood clot;

- difficulty with screw placement can cause injury to the nerves or screw breakage;
- bleeding. Tablets used to 'thin the blood' such as warfarin, aspirin, rivaroxiban and clopidogrel increase the risks of bleeding. You **must** inform your consultant if you are taking these as it is likely you will need to stop taking them before your surgery. Taking medication like non steroidal anti-inflammatories (NSAIDs) could also increase your risk of bleeding and your surgeon will advise you if you need to stop taking these in advance of your operation. If your operation is scheduled with only a week's notice, please check with your consultant or nurse what drugs need to be stopped to prevent your surgery being delayed;
- bone graft non-union or lack of solid fusion (pseudoarthrosis). This can occur in up to 5 out of 100 cases. See page 14 for factors which can affect fusion;
- cage/implant movement can occur in up to 2 out of 100 cases, with 1 out of 100 requiring an operation. In extremely rare cases, cage movement can cause severe damage and cauda equina syndrome (paralysis, bladder or bowel incontinence);
- although rare, the surgery may make your symptoms worse than before.

There are also very rare but serious complications that in extreme circumstances might include:

- damage to the cauda equina and paralysis (the loss of use of the legs, loss of sensation and loss of control of the bladder and bowel. This can occur through bleeding into the spinal canal after surgery (a haematoma). If an event of this nature were to occur, every effort would be made to reverse the situation by returning to the operating theatre to wash out the haematoma. Sometimes however, paralysis can occur as a result of damage or reduction of the blood supply to the nerves or spinal cord and this is, unfortunately, not reversible;
- a stroke, heart attack or other medical or anaesthetic problem; and
- extremely rarely, death; as a result of damage to major blood vessels or vital organs at the front of the spine, which is reported as happening in 1 out of 10,000 cases; or general anaesthetic fatal complications which have been reported in 1 out of 250,000 cases.

ALIF-specific risks and complications

These can include:

- **Damage to the large blood vessels which may result in excessive blood loss.** This is reported as happening in up to 15 out of 100 cases, although it is less common in the hands of an experienced spinal or vascular surgeon. Usually small tears in the vessels can be controlled reasonably simply, though there remains the very small risk of catastrophic bleeding that could, in extremely rare circumstances, lead to death.
- **Sympathetic nerve damage.** There are small nerves directly over the disc space which can be damaged during surgery. These nerves are responsible for many involuntary organ functions, including the heart rate, peristalsis (gut movement), kidney function and, in men, the ability to ejaculate. If these nerves are damaged it can cause problems including:
 - **Retrograde ejaculation (men only).** This is a condition where the valve that causes the ejaculate to be expelled outward during intercourse does not work and the ejaculate takes the path of least resistance, which is up into the bladder. The sensation remains largely the same and this condition does not cause impotence

(the inability to have an erection) but it can unfortunately make conception very difficult. This is reported in fewer than 1 in 100 cases and can resolve over time (a few months to a year).

- o **Warm leg.** This sensation is felt in just one leg, the same side as the surgery has been performed. This can resolve over time but may be a permanent sensation.
- o **Paralytic ileus.** This is a condition where there is an interruption of the normal bowel contraction and the bowel temporarily 'goes to sleep'. It can be a common side effect of abdominal surgery or nerve damage in this type of surgery. Symptoms include constipation and bloating and occasionally vomiting. Diagnosis can be confirmed by a doctor listening to the abdomen with a stethoscope and hearing no bowel sounds. Food should be avoided until sounds are heard and flatus (gas) passed again. This condition can occur in 11 out of 100 cases.

Factors which may affect spinal fusion and your recovery

There are a number of factors that can negatively impact on a solid fusion following surgery, including:

- smoking;
- diabetes or chronic illnesses;
- obesity;
- malnutrition;
- osteoporosis;
- post-surgery activities (see section on recreational activities on page 16); and
- long-term (chronic) steroid use.

Of all these factors, the one that can compromise fusion rate the most is smoking. Nicotine has been shown to be a bone toxin and inhibit the ability of the bone-growing cells in the body (osteoblasts) to grow bone. Patients should make a concerted effort to allow their body the best chance for their bone to heal by not smoking, ideally 2–3 months before the operation. Your surgery may be delayed if you have not stopped smoking (or taking nicotine in another form) beforehand.

What to expect after surgery

Immediately after the operation you will be taken on your bed to the recovery ward where nurses will regularly monitor your blood pressure and pulse. Oxygen will be given to you through a facemask for a period of time to help you recover from the anaesthetic. You will have an intravenous drip until you are able to drink adequately.

A drain (tube) may be placed near the surgical incision if there has been significant bleeding during the operation. This prevents any excess blood or fluid collecting there. The drain will be removed when the drainage has stopped, usually the next day, after surgery.

It is very normal to experience some level of discomfort or back and leg pain after the surgery. The nursing and medical staff will help you to control this with appropriate medication. The symptoms in your legs may fluctuate due to increased swelling around the nerves.

As the nerve becomes less irritated and swollen, your leg pain should settle. This can take eight weeks, or longer. Normal feeling and strength in your legs is likely to take a lot longer and is likely to improve slowly over the next year or so. It is important not to suddenly stop taking certain pain relief medication such as morphine or neuropathic medication (gabapentin, pregabalin or amitriptyline). It will be necessary to gradually 'wean' yourself off them – your GP can advise you if necessary.

The ward physiotherapist will visit you after the operation to teach you exercises and help you out of bed. They will show you the correct way to move safely. Once you are confident and independently mobile, you will be encouraged to practise climbing stairs with the physiotherapist. Once stable you will be allowed home, usually 1–2 days after surgery.

Please arrange for a friend or relative to collect you, as driving yourself or taking public transport is not advised in the initial stages of recovery. If you qualify for patient transport and are likely to require this service, please let one of the nurses know as soon as you can as this may need to be pre-arranged or your discharge home could be delayed.

Wound care

Skin wound closure depends on your surgeon's preference, and include absorbable sutures (stitches), removable sutures or clips (surgical staples).

If you have removable sutures or clips, you will be advised by the ward nurse to arrange an appointment with your GP's practice nurse usually 10 – 14 days after the operation for them to be removed.

If you have absorbable sutures, you will be advised by the ward nurse whether you need to make an appointment with your GP's practice nurse to have a wound check or when you can take off the dressing yourself.

You may shower 48 hours after surgery if you are careful but you must avoid getting the dressing too wet. Most dressings used are 'splash-proof', but if water gets underneath, it will need to be changed. A simple, dry dressing from a pharmacy is sufficient to use. Bathing should be avoided for two weeks.

Please contact your hospital or your GP if you think your wound might be infected. Symptoms could include:

- redness around the wound;
- wound leakage; or
- you have a high temperature.

Once the wound has been checked and if the scar is sensitive to touch, you can start to massage around the scar using a non-perfumed cream or oil to encourage normal sensation and healing.

Driving

Normally you will be advised to avoid driving for four weeks depending on your recovery. If you have no altered sensation or weakness in your legs at that point, then you may resume driving if you feel safe to do so but you must be confident to do an emergency stop. It is advisable not to travel for long distances initially (no longer than 20 minutes), without taking a break to 'stretch your legs'. Gradually increase your sitting tolerance over 4–8 weeks.

Recreational activities

It is important to keep mobile after surgery. You will find you get stiff if sitting for longer than about 20 minutes, so get up and walk about regularly. Walking outside is fine but again, increase your walking distances gradually. You will be advised to avoid lifting anything heavy, certainly for the first few weeks but maybe as long as three months after your operation.

Please check with your consultant and physiotherapist when you are able to resume specific activities, such as swimming or golf, as the advice could range from between six weeks to three months. A graduated return to sport is then advisable. Your surgeon may advise you to avoid flying for six weeks (and long-haul flights for up to three months) because of the increased risk of deep vein thrombosis after surgery.

Work

Returning to work is dependent on both your recovery and your job. Most people are off work for an initial four weeks but if you are in a strenuous job you may need up to eight weeks. It is always sensible to discuss with your employer if you can return on 'light duties' or reduced hours at first. There is usually nothing to stop you doing computer/office work at an earlier date provided you can keep moving about. The hospital will issue you with a fitness to work (off work) certificate or you may ask your GP.

Follow-up

Your surgeon will advise you when you should attend clinic after your operation. If you have any queries about the information in this booklet, please discuss them with the ward nurses or a member of your consultant's team.



What is the British Spine Registry (BSR)?

The British Spine Registry aims to collect information about spinal surgery across the UK. This will help us to find out which spinal operations are the most effective and in which patients they work best. This should improve patient care in the future.

The Registry will enable patient outcomes to be assessed using questionnaires. These will allow surgeons to see how much improvement there has been from treatment.

This has worked for hip and knee joint replacements through the National Joint Registry. We need your help to improve spinal surgery in the UK.

What data is collected?

Your personal details allow the BSR to link you to the surgery you have had. They also allow us to link together all the questionnaires you complete. If you need any further spinal surgery in the future, details of previous operations will be available to your surgeon.

Personal details needed by the BSR are your name, gender, date of birth, address, email address and NHS number.

Your personal details are treated as confidential at all times and will be kept secure. This data is controlled by the British Association of Spine Surgeons (BASS) and held outside the NHS. Personal details will be removed before any data analysis is performed, retaining only age and gender. Your personal data and email address will not be available to anyone outside BASS and its secure IT provider. Anonymised data may be released to approved organisations for approved purposes, but a signed agreement will restrict what they can do with the data so patient confidentiality is protected.

Your personal data is very important, as this will allow us to link details of your diagnosis and surgery with any problems or complications after surgery. You may also be asked to complete questionnaires before and after surgery to work out how successful the surgery has been. This will only be possible if we can connect you to the questionnaires through your personal details.

Do I have to give consent?

No, your participation in the BSR is voluntary and whether you consent or not, your medical care will be the same. Your personal details cannot be kept without your consent. This will be obtained either by asking you to physically sign a consent form or electronically sign one through an email link to a questionnaire or at a questionnaire kiosk in the outpatient clinic.

You can withdraw your consent at any time or request access to your data by:

- going to the patient section of the BSR website at www.britishspineregistry.com; or
- writing to us at the BSR centre (see address on next page). Please state if you are happy for us to keep existing data, but do not want to be contacted, or whether you want your data to be anonymised (so it cannot be identified).

Research

Your consent will allow the BSR to examine details of your diagnosis, surgical procedure, any complications, your outcome after surgery and your questionnaires. These are known as 'service evaluations' or 'audits'.

Operation and patient information, including questionnaires in the BSR, may be used for medical research. The purpose of this research is to improve our understanding and treatment of spinal problems. The majority of our research uses only anonymised information which means it is impossible to identify individuals. From time to time, researchers may wish to gather additional information. In these cases we would seek your approval before disclosing your contact details.

You do not have to take part in any research study you are invited to take part in and saying no does not affect the care you receive.

All studies using data from the Registry will be recorded on the BSR website at www.britishspineregistry.com

Children

Parents are asked to consent for data to be collected from their child. Looking at the outcome of spinal surgical procedures is just as vital in children as it is in adults.

Further information

The BSR website at www.britishspineregistry.com contains more information, including details of any studies and any information obtained through the Registry data.

To contact the BSR, write to:

The British Spine Registry
Amplitude Clinical Services
2nd Floor Orchard House
Victoria Square
Droitwich
Worcestershire WR9 8QT

Produced, researched and revised by spinal nurse specialist Helen Vernau on behalf of the BASS Consent and Patient Information Committee.

Designed and illustrated by Design Services at East Suffolk and North Essex NHS Foundation Trust.